

## AMENDMENTS TO THE CLAIMS


Kindly amend the claims as follows:

In the claims:

Please amend the claims as follows:

---

1. (Previously Amended): A method for inhibiting ALDH-2 in a human comprising contacting ALDH-2 with a compound of formula I



Formula I

wherein:

R is substituted or unsubstituted and is a

sugar moiety;

peptide;

polyether;


straight chain alkyl having 1-11 carbon atoms, or branched chain alkyl having 1-30 carbon atoms, where the branched chain alkyl comprises a straight chain alkyl portion having 1-11 carbon atoms substituted with straight or branched chain lower alkyl groups having 1-6 carbon atoms;

hydroxyalkyl where the alkyl portion is straight chain alkyl having 2-11 carbon atoms, or branched chain alkyl having 2-30 carbon atoms, where the branched chain alkyl comprises a straight

chain alkyl portion having 2-11 carbon atoms substituted with straight or branched chain lower alkyl groups having 1-6 carbon atoms;

aminoalkyl where the alkyl portion is straight chain alkyl having 2-11 carbon atoms, or branched chain alkyl having 2-30 carbon atoms, where the branched chain alkyl comprises a straight chain alkyl portion having 2-11 carbon atoms substituted with straight or branched chain lower alkyl groups having 1-6 carbon atoms;

carboxyalkyl where the alkyl portion is straight chain alkyl having 2-11 carbon atoms, or branched chain alkyl having 2-30 carbon atoms, where the branched chain alkyl comprises a straight chain alkyl portion having 2-11 carbon atoms substituted with straight or branched chain lower alkyl groups having 1-6 carbon atoms; or




where X is straight chain alkylene having 2-11 carbon atoms, or branched chain alkylene having 2-30 carbon atoms, where the branched chain alkylene comprise a straight chain alkylene portion having 2-11 carbon atoms substituted with straight or branched chain lower alkyl groups having 1-6 carbon atoms; and

R' is straight or branched alkyl having 1-6 carbon atoms, in an amount effective to increase concentration of 5-hydroxyindole-3-acetaldehyde or 3,4-dihydroxyphenylacetaldehyde.

2. (Presently Amended): The method of claim 1 wherein the sugar moiety is glucosyl, L or D aldo or keto-tetrose, pentose, heptose, an amino, alcohol or acid derivative of tetrose, pentose, hexose or heptose, a deoxy-analog of tetrose, pentose, hexose or heptose.

3. (Presently Amended): A method for therapeutically reducing alcohol consumption in a

human in need thereof comprising administering to the human a compound of formula I



Formula I

wherein:

*10' cont*

R is substituted or unsubstituted and is a sugar moiety selected from the group consisting of L or and D, aldo- or and keto-, tetroses, pentoses, ~~hexoses~~, heptoses, amino derivatives of tetroses, pentoses, or heptoses, alcohol derivatives of tetroses, pentoses, or heptoses, and/or acid derivatives of tetroses, pentoses, or heptoses and deoxy analogs of tetroses, pentoses, or heptoses thereof;

peptide;

polyether; or

aminoalkyl where the alkyl portion is straight chain alkyl having 2-11 carbon atoms, or branched chain alkyl having 2-30 carbon atoms, where the branched chain alkyl comprises a straight chain alkyl portion having 2-11 carbon atoms substituted with straight or branched chain lower alkyl groups having 1-6 carbon atoms;

in an amount effective to increase concentration of an biogenic aldehyde formed during catabolism of a neurotransmitter.


4. (Cancelled)

5. (Original): The method of claim 3 wherein the neurotransmitter is serotonin or dopamine.

6. (Previously Amended): The method of claim 3 wherein said biogenic aldehyde is 5-hydroxyindole-3-acetaldehyde or 3,4-dihydroxyphenyl-3-acetaldehyde.

7-12 (Cancelled)

13. (New): A method for therapeutically reducing alcohol consumption in a human in need thereof comprising administering to the human a compound of formula I



Formula I

wherein:

R is substituted or unsubstituted and is a

peptide;

polyether; or

aminoalkyl where the alkyl portion is straight chain alkyl having 2-11 carbon atoms, or branched chain alkyl having 2-30 carbon atoms, where the branched chain alkyl comprises a straight chain alkyl portion having 2-11 carbon atoms substituted with straight or branched chain lower alkyl

groups having 1-6 carbon atoms;

in an amount effective to increase concentration of an aldehyde formed during catabolism of a neurotransmitter.

20' cont  
14. (New): The method of claim 13 wherein the neurotransmitter is serotonin or dopamine.

15. (New): The method of claim 13 wherein the aldehyde is 5-hydroxyindole-3-acetaldehyde or 3,4-dihydroxyphenyl-3-acetaldehyde.

---